



TIMM8A gene

translocase of inner mitochondrial membrane 8A

Normal Function

The *TIMM8A* gene provides instructions for making a protein that is found inside mitochondria, which are structures within cells that convert the energy from food into a form that cells can use. Mitochondria have two membranes, an outer membrane and an inner membrane, which are separated by a fluid-filled area called the intermembrane space. The TIMM8A protein is found in the intermembrane space, where it forms a complex (a group of proteins that work together) with a very similar protein called TIMM13. This complex transports other proteins across the intermembrane space to the mitochondrial inner membrane.

Health Conditions Related to Genetic Changes

deafness-dystonia-optic neuropathy syndrome


At least 15 mutations in the *TIMM8A* gene have been found to cause deafness-dystonia-optic neuropathy (DDON) syndrome. Most of these mutations result in the absence of functional TIMM8A protein inside the mitochondria, which prevents the formation of the TIMM8A/TIMM13 complex. Researchers believe that the lack of this complex leads to abnormal transport of proteins across the intermembrane space, although it is unclear how abnormal protein transport affects the function of the mitochondria and causes the signs and symptoms of DDON syndrome.

Some people with DDON syndrome have large DNA deletions that remove the entire *TIMM8A* gene and one end of a neighboring gene known as *BTK*. Mutations in the *BTK* gene cause X-linked agammaglobulinemia (XLA), which is characterized by an increased susceptibility to infections. Individuals with large DNA deletions that include the *TIMM8A* gene and the *BTK* gene have the signs and symptoms of both DDON syndrome and XLA.

Chromosomal Location

Cytogenetic Location: Xq22.1, which is the long (q) arm of the X chromosome at position 22.1

Molecular Location: base pairs 101,345,656 to 101,348,969 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- DDP
- DDP1
- deafness/dystonia peptide
- DFN1
- MGC12262
- TIM8A_HUMAN
- translocase of inner mitochondrial membrane 8 homolog A
- translocase of inner mitochondrial membrane 8 homolog A (yeast)

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): The Mitochondrion
<https://www.ncbi.nlm.nih.gov/books/NBK26894/>

GeneReviews

- Deafness-Dystonia-Optic Neuronopathy Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1216>

Scientific Articles on PubMed

- PubMed

<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28TIMM8A%5BTIAB%5D%29+OR+%28translocase+of+inner+mitochondrial+membrane+8+homolog+A%5BTIAB%5D%29%29+OR+%28%28DDP%5BTIAB%5D%29+OR+%28DDP1%5BTIAB%5D%29+OR+%28DFN1%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+NOT+%28%28cancer%29+OR+%28tumor%29+OR+%28anticancer%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>

OMIM

- TRANSLOCASE OF INNER MITOCHONDRIAL MEMBRANE 8, YEAST, HOMOLOG OF, A
<http://omim.org/entry/300356>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_TIMM8A.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=TIMM8A%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=11817
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/1678>
- UniProt
<http://www.uniprot.org/uniprot/O60220>

Sources for This Summary

- Binder J, Hofmann S, Kreisel S, Wöhrle JC, Bätzner H, Krauss JK, Hennerici MG, Bauer MF. Clinical and molecular findings in a patient with a novel mutation in the deafness-dystonia peptide (DDP1) gene. *Brain.* 2003 Aug;126(Pt 8):1814-20. Epub 2003 Jun 4. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12805099>
- GeneReview: Deafness-Dystonia-Optic Neuronopathy Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1216>
- Jyonouchi H, Geng L, Törnér GA, Vinekar K, Feng D, Fitzgerald-Bocarsly P. Monozygous twins with a microdeletion syndrome involving BTK, DDP1, and two other genes; evidence of intact dendritic cell development and TLR responses. *Eur J Pediatr.* 2008 Mar;167(3):317-21. Epub 2007 May 23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17520285>

- Richter D, Conley ME, Rohrer J, Myers LA, Zahradka K, Kelecic J, Sertic J, Stavljenic-Rukavina A. A contiguous deletion syndrome of X-linked agammaglobulinemia and sensorineural deafness. *Pediatr Allergy Immunol.* 2001 Apr;12(2):107-11.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11338284>
- Roesch K, Curran SP, Tranebaerg L, Koehler CM. Human deafness dystonia syndrome is caused by a defect in assembly of the DDP1/TIMM8a-TIMM13 complex. *Hum Mol Genet.* 2002 Mar 1;11(5):477-86.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11875042>
- Roesch K, Hynds PJ, Varga R, Tranebaerg L, Koehler CM. The calcium-binding aspartate/glutamate carriers, citrin and aralar1, are new substrates for the DDP1/TIMM8a-TIMM13 complex. *Hum Mol Genet.* 2004 Sep 15;13(18):2101-11. Epub 2004 Jul 14.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15254020>
- Sedivá A, Smith CI, Asplund AC, Hadac J, Janda A, Zeman J, Hansíková H, Dvoráková L, Mrázová L, Velbri S, Koehler C, Roesch K, Sullivan KE, Futatani T, Ochs HD. Contiguous X-chromosome deletion syndrome encompassing the BTK, TIMM8A, TAF7L, and DRP2 genes. *J Clin Immunol.* 2007 Nov;27(6):640-6. Epub 2007 Sep 12.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17851739>
- OMIM: TRANSLOCASE OF INNER MITOCHONDRIAL MEMBRANE 8, YEAST, HOMOLOG OF, A
<http://omim.org/entry/300356>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/TIMM8A>

Reviewed: September 2008

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services